

Review article

Traditional Methods of Diabetes Mellitus Treatment

Khalid Abdulkareem Mohammed^{1*}

ABSTRACT

Diabetes mellitus is one of the most common metabolic diseases in the world, and the incidence of this disease is not limited to specific age groups, and it appears among men and women and there is no effect on gender, age, and geographic area on the outbreak of this disease. Synthetic and natural materials have been used to control blood sugar levels in diabetics. The current study sheds light on the importance of this disease and the extent of its spread in the world, in addition to its most important types, which are Insulin-dependent diabetes mellitus (IDDM) Insulin-independent diabetes mellitus (IIDM), and Gestational diabetes. The current study showed the most important herbal extracts that were used and are still used today in many countries to control hyperglycemia in diabetics, especially *Brassica oleracea*. The effect of these herbal extracts is due to the presence of some active substances that reduce high blood sugar levels, and the most important of these substances are phenols and alkaloids. The effect of these substances on insulin-producing cells was discussed, as well as their direct and indirect effect on controlling high blood sugar in laboratory animals and people suffering from various types of diabetes. The study concluded that the use of herbal medicine is effective in reducing hyperglycemia in the patients with diabetes, in addition to being safe and having no side effects on patients who use them.

Keywords: Diabetes mellitus, Herbal medicine, Insulin-dependent diabetes, Insulin-independent diabetes.

Citation: Mohammed KA. (2023) Traditional methods of diabetes mellitus treatment. *World J Exp Biosci* 11: 1- 6. <https://doi.org/10.65329/wjeb.v11.01.001>

Received December 2, 2022; Accepted January 21, 2023; Published: February 3, 2023.

1. INTRODUCTION

Most conventional approaches and remedies utilized for the management of diabetes rely on the utilization of herbal medicine. Herbal medicine entails the utilization of plants for medicinal intentions. Throughout a significant portion of human history, plants have constituted the foundation for medical treatments, and this form of traditional medicine continues to be widely practiced in the current era [1]. One of the most practical plants employed in herbal medicine is *Brassica oleracea*. *B. oleracea* pertains to the Brassicaceae (Cruciferae) botanical family. The wild variant of this plant originated along the Atlantic coast of Western Europe and the Mediterranean basin. *B. oleracea* has been cultivated as a vegetable for over 2,500 years, and through selective breeding, distinct characteristics of the plant have been developed [2]. Diabetes encompasses a chronic metabolic disorder typified by elevated blood glucose le-

vels. This condition arises either due to inadequate endogenous insulin production by the pancreatic beta cells, known as type 1 diabetes, or impaired insulin secretion and/or action, referred to as type 2 diabetes. Type 1 diabetes represents an autoimmune ailment characterized by T-cell-mediated destruction of the pancreatic beta cells. In the case of type 2 diabetes, there is a gradual progression of insulin resistance and beta cell dysfunction, which are strongly linked to obesity and a sedentary lifestyle [3]. Given the higher incidence of the risk factors, the prevalence of diabetes is escalating worldwide, with developing nations experiencing a more pronounced impact. Current estimates project a 69% increase in the number of adults affected by this disease between 2010 and 2030, in contrast to a 20% increase observed in developed countries [4]. Significant advancements have been made in the field of synthetic drug development. However, ongoing investigations are being carried out to explore natural and cost-effective food

* Correspondence: Assistant Professor, Dr. Khalid Abdulkareem Mohammed. E. mail: khalid.kreem@yahoo.com

Department Chemistry, College of Science, University of Baghdad, Baghdad, Iraq.

Full list of author information is available at the end of the article.

Copyright: © Khalid A. Mohammed. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY4.0), which permits unrestricted use, distribution, and reproduction in any site, provided the original author and source are credited.

sources for managing hyperglycemia and hypertension associated with diabetes. This is achieved through the consumption of diets rich in legumes, fruits, vegetables, herbs, and spices. These plant-based foods contain essential nutrients such as vitamins, minerals, and dietary fibers, as well as bioactive compounds including polyphenols, carotenoids, and alkaloids [5,6]. These bioactive compounds have specific structure-function benefits [7] and have the potential to prevent and reduce the occurrence of early stages of chronic diseases such as type II diabetes and cardiovascular diseases, along with their complications. Many plant foods and specific varieties of certain species contain hypoglycemic compounds that act as anti-metabolites, effectively blocking specific disease pathways, including the oxidation pathway of fatty acids [5,8]. Medicinal herbs and seeds derived from plant sources are known to have low toxicity levels and possess various therapeutic properties. Consequently, they are considered to be safer alternatives to synthetic drugs. Natural plant and food sources also offer α -amylase and α -glucosidase inhibitors, which contribute to controlling postprandial hyperglycemia [9] with minimal side effects [10].

The levels of phenolic phytochemicals in leaves, stems, fruits, and seeds of numerous plant sources are being evaluated, as they are known to exhibit high antioxidant activity [11]. Phenolic phytochemicals serve as a defense mechanism for plants against abiotic and biotic stresses, but they also offer significant benefits in preventing and combating chronic diseases associated with oxidative stress in humans [12]. The presence of specific phenolic compounds in plant foods is strongly correlated with high antioxidant activities and substantial amounts of α -amylase and α -glucosidase inhibitors. These compounds play a vital role in managing hyperglycemia and related complications of hypertension [10].

The utilization of artificial drugs in the treatment of diabetes and glucose control in diabetic patients is often accompanied by various side effects [13]. Furthermore, these drugs tend to be costly, prompting many research centers worldwide to focus on the use of natural plant-based products for glucose level control in diabetic patients. The present study offers substantial evidence regarding the effectiveness of different compounds extracted from *B. oleracea* var. *capitata* and other plants in reducing blood glucose levels in diabetic patients [14].

2. DIABETES MELLITUS

Diabetes mellitus, commonly known as diabetes, refers to a collection of metabolic diseases characterized by prolonged periods of elevated blood sugar levels. This abnormal increase in blood sugar manifests in symptoms such as frequent urination, increased thirst, and heightened hunger. If left untreated, diabetes can lead to various complications. These complications include acute conditions like diabetic ketoacidosis and nonketotic hyperosmolar coma, as well as serious long-term complications such as heart disease, stroke, kidney failure, foot ulcers, and damage to the eyes [15]. Diabetes mellitus is now emerging as a prevalent metabolic disorder, arising from the inability of the body to properly respond to high blood glucose levels. Type II diabetes mellitus accounts for approximately 90-95% of all diagnosed diabetic cases [16]. The epidemic nature of type II diabetes is closely linked to obesity, with the World Health Organization (WHO) reporting that over 220 million individuals were affected by the condition in 2009 [17]. Projections suggest that the number of people afflicted with diabetes will exceed 350 million by 2030 [18]. The incidence of diabetes patients experiencing complications is steadily

increasing, leading to hospital admissions due to diabetes itself or its associated complications, such as stroke, hypertension, amputation, nephropathy, neuropathy, retino-pathy, cardiovascular issues, impotence, and skin lesions. An analysis of mortality trends in the United States between 1970 and 2002 revealed the largest reductions in age-adjusted death rates were for stroke (63%) and heart disease (52%) [19,20]. The growing number of individuals affected by diabetes indicates a global epidemic. The prevalence of this disease was estimated to be 2.8% in 2000 and is projected to rise to 4.4% by 2030. While the prevalence of diabetes is anticipated to increase across all age groups, it is expected that there will be a particularly pronounced increase in the proportion of patients aged 45-64 years [21]. A previous study demonstrated that individuals with diabetes who have inadequate health literacy and hypertension possess less knowledge about their condition compared to those with sufficient health literacy [22].

2.1. Insulin-dependent diabetes mellitus (IDDM)

Diabetes mellitus results from the body's failure to produce enough insulin. This form was previously referred to as IDDM (type 1 diabetes) or "juvenile diabetes". The cause is unknown. Type 1 diabetes occurs as a result of auto-immune beta-cell destruction in the pancreas, characterized by a total absence of insulin production. Type 1 diabetes is responsible for 5% to 10% of all cases of diabetes. Associated risk factors include autoimmune, genetic, and environmental factors. Till the present time, known solutions to prevent diabetes have not been discovered [23]. Type 1 diabetes onset is relatively early in life, in childhood or adolescence, and usually before the age of thirty. This type of diabetes is a relatively homogeneous disease in which the insulin secretion of beta cells in the pancreas declines and eventually ceases totally [24, 25].

2.2. Insulin-independent diabetes mellitus (IIDM)

Type 2 diabetes can be linked to be accounting for around 90 percent of all cases, it is a chronic metabolic disorder, in which the body is unable to utilize glucose from food because of the inability of the pancreas to produce insulin or produces insufficient insulin, or the insulin itself is inactive [26]. Type 2 develops when there is an unexpected increased resistance against the action of insulin and the body cannot produce proportionate insulin to counter the resistance. The incidence of Type 2 diabetes in children and adolescents is noted to be on a dramatic increase. It accounts for 90% to 95% of all diagnosed cases of diabetes [23, 27].

2.3. Gestational diabetes

Gestational diabetes is the third main case and occurs when pregnant women without a previous history of diabetes develop a high blood glucose level.

3. DIANETES MANAGEMENT BY TRADITIONAL HERBAL MEDICINES

A multitude of scholarly articles have been published regarding the use of medicinal plants in the treatment of diabetes in various regions across the globe [28]. These articles have also explored the specific use of medicinal plants in certain areas, such as West Africa [29], Central America [30], and Asia [31].

These reviews have shed light on the fact that a significant portion of the global population relies on traditional medicine for the management of diabetes. This reliance on traditional medicine has been acknowledged by the World Health Organization (WHO), which estimates that approximately 80% of the populations in African and Asian countries depend on traditional medicine for their primary healthcare needs [32]. The WHO recognizes traditional medicine as an accessible, affordable, and culturally accepted form of healthcare that is trusted by a large number of people. It is seen as a way to address the increasing prevalence of chronic non-communicable diseases, despite rising healthcare costs and global austerity measures [32].

Although significant progress has been made in the development of synthetic drugs, researchers are currently investigating natural and cost-effective food sources for managing hyperglycemia and hypertension in the early stages of diabetes. This research focuses on diets that are rich in legumes, fruits, vegetables, herbs, and spices. These plant-based foods provide essential nutrients, such as vitamins, minerals, and dietary fibers, as well as bioactive compounds like polyphenols and carotenoids [5]. These bioactive compounds have specific structure-function benefits that are beneficial for managing diabetes [33]. Many plant-based foods and specific varieties of certain species contain compounds that have hypoglycemic properties. These compounds act as anti-metabolites, blocking specific disease pathways, including the oxidation pathway of fatty acids [8]. Medicinal herbs and seeds derived from plants are known to have low toxicity levels and possess various therapeutic properties, making them a safer alternative to synthetic drugs. Natural plant and food sources also provide α -amylase and α -glucosidase inhibitors, which can help control postprandial hyperglycemia with minimal side effects [7, 34].

Researchers are currently evaluating the levels of phenolic phytochemicals in herbs and seeds derived from various plant sources. These phytochemicals have high antioxidant activity and are used by plants to protect themselves from environmental and biological stresses. They also have potential benefits in preventing and treating chronic diseases in humans that are linked to oxidative stress [35]. The presence of certain phenolic compounds in plant-based foods is associated with high antioxidant activity, as well as high levels of α -amylase and α -glucosidase inhibitors. These compounds play a role in managing hyperglycemia and related complications, such as hypertension [36].

3.1. Medicinal Plants with Antidiabetic

Many examples of plants and their extract that used in diabetes treatment in several places all over the world there are several examples of plants uses for controlling the blood sugar and managing the debits such as, *Abelmoschus moschatus* Medik (Malvaceae) (Liu et al., 2007), *Acacia arabica* (Lam) Wild. (Mimosaceae), *Achyranthes aspera* L (Amaranthaceae), *Acosmium panamense* Schott. (Leguminosae), *Allium cepa* L. (onion): (Liliaceae), *Allium sativum* L. (garlic): (Liliaceae), *Annona squamosa* L (Annonaceae), *Artemisia herba-alba* Asso (Med). (Asteraceae), *Astragalus membranaceus* Bunge (Fisch.) (Leguminosae), *Bauhinia candicans* Benth (Leguminosae), *Bauhinia forficata* Link. (Caesalpiniaceae), *Bidens pilosa* L (Asteraceae), *Biophytum sensitivum* (L) DC. (Oxalidaceae), *Bixa orellana* L. (Bixaceae), *Brassica nigra* (L) Koch (Brassicaceae), *Bryonia alba* L. (Cucurbitaceae), *Caesalpinia bonducuella* (L) Roxb. (Caesalpiniaceae), *Cajanus*

cajan (L) Millsp. (Papilionaceae), *Carum carvi* L (CC) (Apiaceae) / *Capparis spinosa* L. (CS) (Capparidaceae), *Casearia esculenta* Roxb. (Flacourtiaceae), *Cassia auriculata* L. (Caesalpiniaceae), *Catharanthus roseus* (L) G. Don. (Apocynaceae), *Chamaemelum nobile* (L) All. (Asteraceae), *Cichorium intybus* L. (Asteraceae), *Clausena anisata* (Willd) Benth. (Rutaceae), *Coriandrum sativum* L (Apiaceae), *Cuminum cyminum* L (Apiaceae), *Cuminum nigrum* L (Apiaceae), *Cyamopsis tetragonoloba* (L) Taubert. (Papilionaceae), *Dioscorea dumetorum* (Kunth) Pax. (Dioscoreaceae), *Ericostema littorale* blume (Gentianaceae), *Fraxinus excelsior* L (Oleaceae), *Garcinia kola* Heckel (W & C Afr) (Clusiaceae), *Gongronema latifolium* Endl. (Asclepiadaceae), *Helicteres isora* L., As. (Sterculiaceae), *Hypoxis hemerocallidea* conn Corm (African potato) (Hypoxidaceae), *Inula racemosa* Hook.f. (Asteraceae) *Lepidium sativum* L. (Brassicaceae), *Mangifera indica* L. (Anacardiaceae), *Momordica charantia* L. (Cucurbitaceae), *Morinda lucida* Benth. (Rubiaceae), *Myrcia uniflora* Barb., Rods. (Myricaceae), *Ocimum sanctum* L. (Lamiaceae), *Pterocarpus marsupium* Roxb. (Papilionaceae), *Retama raetam* (RR) (Forsk) Webb. (Papilionaceae), *Salacia reticulata* W. (Celastraceae), *Telfaria occidentalis* Hook. (Cucurbitaceae) [37], and *Brassica oleracea* var. *capitata*. There are several components that participate in controlling the sugar in the patients of diabetes such as phenolic and alkaloid compounds [14].

3.2. Phenolic compounds

A diverse array of phenolic compounds have been identified as the active components in some of the plants discussed in this review. For instance, anthraquinone glycosides derived from *Morinda citrifolia* exhibited a reduction in blood glucose levels in STZ-induced diabetic mice at a dosage of 100 mg/kg [38], which also can be clear in the case of *B. oleracea* var. *capitata* [14]. Despite this, the use of a registered herbal product called Tahitian noni juices (TNJ) is quite prevalent in Nigeria for various ailments, including diabetes. The administration of 1ml / 150 mg body weight of TNJ to rats twice daily for four weeks, both before and after the induction of diabetes with alloxan, resulted in a significant decrease in blood sugar levels, suggesting a preventive effect of the extract against alloxan-induced diabetes [39]. However, the presence of these phenolic compounds in the marketed product has not been confirmed. Kolaviron, a mixture of flavanones isolated from *Garcinia kola* Heckel (bitter kola), which is highly regarded in many parts of West Africa, has been found to lower blood sugar levels in normal and alloxan-induced diabetic mice at a dosage of 100 mg/kg, as well as inhibit rat lens aldose reductase (RLAR) activity [40]. Furthermore, several isolated flavonoids have been identified as bioactive constituents. Isoscutellarein (8-hydroxyapigenin), a flavonoid isolated from the hot water extract of *Bixa orellana* L. leaves, has been identified as an aldose reductase inhibitor. Rutin and quercetin, isolated from *Bauhinia monandra* Kurz leaves, have been identified as anti-hyperglycemic constituents in alloxan-induced diabetic rats [41]. A bioassay-guided fractionation of the stem bark of *Cassia fistula* L. led to the identification of catechin as the bioactive agent. Catechin was found to decrease plasma glucose levels in STZ-induced diabetic rats, with direct effects on glucose metabolizing enzymes and the expression of the glucose transporter GLUT4 [42].

The antioxidant properties of flavonoids, particularly their ability to scavenge free radicals, have been associated with the presence of aromatic hydroxyl groups in the benzo- γ -pyran

structure. These properties have been shown to protect pancreatic islet cells from oxidative stress and aid in the regeneration of β -cells, as demonstrated by epicatechin found in green tea [43] and quercetin [44]. More importantly, these flavonoids can prevent the formation of advanced glycated end products (AGEs) and other diabetic complications that arise from high levels of oxidative stress, such as atherosclerosis, nephropathy, neuropathy, retinopathy, and erectile dysfunction [45]. Therefore, the presence of quercetin, epicatechin, and other potent antioxidant flavonoids in a wide range of plants, including *Irvingia gabonensis* and *B. oleracea* var. *capitata*, contributes to their use in the holistic management of diabetes and the prevention of diabetic complications [14].

Other types of flavonoids have also demonstrated a direct impact on specific therapeutic targets in the context of diabetes. To illustrate, the introduction of naringin or hesperidin into the diet of mice has been observed to modulate the activity of enzymes responsible for glucose metabolism. Specifically, there was an increase in hepatic glucokinase activity and a decrease in hepatic glucose-6-phosphatase activity in diabetic mice [46] and GK type-2 diabetic rats [47]. These two particular flavonoids are present in all citrus fruits and have also been identified in *Senna alata* [48] and *Rauvolfiavomitoria* [49], suggesting they may contribute to the observed effects. Myricetin is another flavonoid that has demonstrated direct positive effects in the realm of diabetes, specifically through enhanced glycogen metabolism and improved insulin sensitivity [50].

3.3. Alkaloids

A number of investigations have assessed the antidiabetic properties of alkaloids derived from various plant components [51]. Odoh and Ezugwu (2012) carried out an examination to establish the effect of alkaloids from *Acanthus montanus* leaf (AAML) on diabetes. The experiment involved the intraperitoneal administration of AAML to alloxan-induced diabetic rats at doses of 100, 200, and 400mg/kg for durations of 4 and 8 weeks. The researchers discovered that the intraperitoneal administration of AAML alkaloids was safe and could potentially provide beneficial effects as an immunostimulant, hepatoprotective, and hypocholesterolemic agent when administered over an extended period of time [52]. Tiong et al. (2013) documented the antioxidant and antidiabetic activities of the primary alkaloids isolated from the extract of *C. roseus* (L.) G. Don leaves [53]. Agrawal et al. (2013) revealed that the methanol extract of *Aerva lanata* Linn. roots (AL) was fractionated using different solvents. The partially purified alkaloid basified toluene fraction (PPABTF) at a dose of 20mg/kg exhibited significant effects on streptozotocin-nicotinamide-induced type-II NIDDM in rats. This activity may be attributed to the presence of alkaloids such as canthin-6-one derivatives. Therefore, numerous investigations have demonstrated the antidiabetic effects of alkaloids derived from diverse plants [54]. Most studies conducted by various researchers have focused on the direct impact of phenolic compounds and alkaloids derived from different plant parts on reducing glucose levels in experimental animals, including rats, rabbits, and mice [55].

4. CONCLUSION

The current study showed that the use of various herbs to control of sugar in blood of patients suffering with diabetes is safe and free of many of the side effects that can appear from the use of chemically manufactured drugs. The study showed that there are large numbers of herbs that can be used to control

diabetes. Phenol and alkaloids play an important role in maintaining blood sugar levels in diabetic' patients.

Acknowledgments

I would like to take this opportunity to extend my sincere thanks to all members of the Department of Chemistry at the University of Baghdad as well as to the head of the department for the support that was provided to me during the period of preparing this article.

Funding information

This work received no specific grant from any funding agency.

Conflict of interest

The authors declare that they have no conflict of interests.

Ethical Approval

This review was approved by the Ethical Committee of the University of Baghdad, Baghdad, Iraq (No 214, 2022).

Author contributions

Khalid A. Mohammed. Conceptualization; Data curation; Formal analysis; Investigation; Resources; Supervision; Validation; Visualization; Roles/Writing – original draft; and Writing – review & editing.

5. REFERENCES

- [1] Ghosh SS, Gehr TW, Ghosh S. (2014) Curcumin and chronic kidney disease (CKD): major mode of action through stimulating endogenous intestinal alkaline phosphatase. *Molecules* **19**:20139-56. doi: <https://doi.org/10.3390/molecules191220139>. PMID: 25474287; PMCID: PMC6271001.
- [2] Yousefimehr F, Jafarirad S, Salehi R, Zakerhamidi MS. (2021) Facile fabricating of rGO and Au/rGO nanocomposites using *Brassica oleracea* var. gongylodes biomass for non-invasive approach in cancer therapy. *Sci Rep* **11**:11900. doi: <https://doi.org/10.1038/s41598-021-91352-7>. PMID: 34099785; PMCID: PMC8184758.
- [3] Tan T, Xiang Y, Deng C, Cao C, Ren Z, et al. (2022) Variable frequencies of peripheral T-lymphocyte subsets in the diabetes spectrum from type 1 diabetes through latent autoimmune diabetes in adults (LADA) to type 2 diabetes. *Front Immunol* **13**:974864. doi: <https://doi.org/10.3389/fimmu.2022.974864>. PMID: 36091068; PMCID: PMC9449581.
- [4] Shaw JE, Sicree RA, Zimmet PZ. (2009) Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract* **87**:4-14. doi: <https://doi.org/10.1016/j.diabres.2009.10.007>. Epub 2009 Nov 6. PMID: 19896746.
- [5] Robert L, Narcy A, Rock E, Demigne C, Mazur A, Rémy C. (2006) Entire potato consumption improves lipid metabolism and antioxidant status in cholesterol-fed rat. *Eur J Nutr* **45**:267-74. doi: <https://doi.org/10.1007/s00394-006-0594-y>. PMID: 16586150.
- [6] Sellamuthu PS, Arulselvan P, Fakurazi S, Kandasamy M. (2014) Beneficial effects of mangiferin isolated from *Salacia chinensis* on biochemical and hematological parameters in rats with streptozotocin-induced diabetes. *Pak J Pharm Sci* **27**:161-7. PMID: 24374436.
- [7] Kwon YI, Apostolidis E, Kim YC, Shetty K. (2007) Health benefits of traditional corn, beans, and pumpkin: in vitro studies for hyperglycemia and hypertension management. *J Med Food* **10**:266-75. doi: <https://doi.org/10.1089/jmf.2006.234>. PMID: 17651062.
- [8] Ahmad Aufa Z, Hassan FA, Ismail A, Mohd Yusof BN, Hamid M. (2014) Chemical compositions and antioxidative and antidiabetic properties of underutilized vegetable palm hearts from *Plectocomiopsis geminiflora* and *Eugeissona insignis*. *J Agric Food Chem* **62**:2077-84. doi: <https://doi.org/10.1021/jf403481p>. PMID: 24499380.
- [9] Riyaphan J, Jhong C-H, Lin S-R, Chang C-H, Tsai M-J, et al. (2018) Hypoglycemic Efficacy of Docking Selected Natural Compounds against α -Glucosidase and α -Amylase. *Molecules* **23**:2260. doi: <https://doi.org/10.3390/molecules23092260>.
- [10] Kim GN, Kwon YI, Jang HD. (2011) Mulberry leaf extract reduces postprandial hyperglycemia with few side effects by inhibiting α -glucosidase in normal rats. *J Med Food* **14**:712-7. doi: <https://doi.org/10.1089/jmf.2010.1368>. PMID: 21631361.

[11] Fu LZ, Zhao LM, Lyu HQ, Yan MQ, Zheng YQ, et al. (2019) Effects of nitrogen level on growth of *Tetrastigma hemsleyanum* and phytochemical content and antioxidant activity in stems and leaves. *Zhongguo Zhong Yao Za Zhi* **44**:696-702. Chinese. doi: <https://doi.org/10.19540/j.cnki.cjcm.20181204.006>. PMID: 3098881.

[12] Guan R, Van Le Q, Yang H, Zhang D, Gu H, et al. (2021) A review of dietary phytochemicals and their relation to oxidative stress and human diseases. *Chemosphere* **271**:129499. doi: <https://doi.org/10.1016/j.chemosphere.2020.129499>. Epub 2021 Jan 4. PMID: 33445014.

[13] Iizuka K. (2022) Is the Use of Artificial Sweeteners Beneficial for Patients with Diabetes Mellitus? The Advantages and Disadvantages of Artificial Sweeteners. *Nutrients* **14**:4446. doi: <https://doi.org/10.3390/nu14214446>.

[14] Uuh-Narvaez JJ, Segura-Campos MR. (2021) Cabbage (*Brassica oleracea* var. *capitata*): A food with functional properties aimed to type 2 diabetes prevention and management. *J Food Sci* **86**:4775-4798. doi: 10.1111/1750-3841.15939. Epub 2021 Oct 17. PMID: 34658044. <https://doi.org/10.1111/1750-3841.15939>.

[15] Lin KD, Hsu CC, Ou HY, Wang CY, Chin MC, Shin SJ. (2019) Diabetes-related kidney, eye, and foot disease in Taiwan: An analysis of nationwide data from 2005 to 2014. *J Formos Med Assoc* **118** Suppl:S103-S110. doi: <https://doi.org/10.1016/j.jfma.2019.07.027>. Epub 2019 Aug 30. PMID: 31477486.

[16] O'Brien JA, Patrick AR, Caro J. (2000) Estimates of direct medical costs for microvascular and macrovascular complications resulting from type 2 diabetes mellitus in the United States in 2000. *Clin Ther* **25**:1017-38. doi: [https://doi.org/10.1016/s0149-2918\(03\)80122-4](https://doi.org/10.1016/s0149-2918(03)80122-4). PMID: 12852716.

[17] Liang B, Wu XY. (2011) Progress on nonhuman primate models of diabetes mellitus. *Dongwuxue Yanjiu* **32**:91-6. Chinese. doi: <https://doi.org/10.3724/SP.J.1141.2011.01091>. PMID: 21341390.

[18] Suija K, Kivistö K, Sarria-Santamera A, Kokko S, Liseckiene I, et al. (2015) Challenges of audit of care on clinical quality indicators for hypertension and type 2 diabetes across four European countries. *Fam Pract* **32**:69-74. doi: <https://doi.org/10.1093/fampra/cmu078>. Epub 2014 Nov 18. PMID: 25411423.

[19] Chaturvedi N. (2007) The burden of diabetes and its complications: trends and implications for intervention. *Diabetes Res Clin Pract* **76** Suppl 1:S3-12. doi: <https://doi.org/10.1016/j.diabres.2007.01.019>. Epub 2007 Mar 6. PMID: 17343954.

[20] Parissis JT, Rafouli-Stergiou P, Mebazaa A, Ikonomidis I, Bistola V, et al. (2012) Acute heart failure in patients with diabetes mellitus: clinical characteristics and predictors of in-hospital mortality. *Int J Cardiol* **157**:108-13. doi: <https://doi.org/10.1016/j.ijcard.2011.11.098>. Epub 2011 Dec 15. PMID: 22178056.

[21] Adiseshiah M. (2005) Effective care of patients with type 2 diabetes and dyslipidemia: a nurse's perspective. *Diabetes Res Clin Pract* **68**:S23-7. doi: <https://doi.org/10.1016/j.diabres.2005.03.017>. Epub 2005 Apr 14. PMID: 15953503.

[22] Wang B, Song WH, Liu GZ. (2005) The effect long-term administration of a selective beta1 blocker bisoprolol on glucose metabolism in patients with essential hypertensive and type 2 diabetes mellitus. *Zhonghua Nei Ke Za Zhi* **44**:503-5. Chinese. PMID: 16080839.

[23] Deshpande AD, Harris-Hayes M, Schootman M. (2008) Epidemiology of diabetes and diabetes-related complications. *Phys Ther* **88**:1254-64. doi: <https://doi.org/10.2522/pti.20080020>. Epub 2008 Sep 18. PMID: 18801858; PMCID: PMC3870323.

[24] Fousteri G, Ippolito E, Ahmed R, Hamad ARA. (2017) Beta-cell Specific Autoantibodies: Are they Just an Indicator of Type 1 Diabetes? *Curr Diabetes Rev* **13**:322-329. doi: <https://doi.org/10.2174/1573399812666160427104157>. PMID: 27117244; PMCID: PMC5266674.

[25] Potter KJ, Westwell-Roper CY, Klimek-Abercrombie AM, Warnock GL, Verchere CB. (2014) Death and dysfunction of transplanted β -cells: lessons learned from type 2 diabetes? *Diabetes* **63**:12-9. doi: <https://doi.org/10.2337/db12-0364>. PMID: 24357689.

[26] Naemiratch B, Manderson L. (2007) Lay Explanations of Type 2 Diabetes in Bangkok, Thailand. *Anthropol Med* **14**:83-94. doi: <https://doi.org/10.1080/13648470601106418>. PMID: 26873802.

[27] Dabelea D, Stafford JM, Mayer-Davis EJ, D'Agostino R Jr, Dolan L, et al. (2017) SEARCH for Diabetes in Youth Research Group. Association of Type 1 Diabetes vs Type 2 Diabetes Diagnosed During Childhood and Adolescence With Complications During Teenage Years and Young Adulthood. *JAMA* **317**:825-835. doi: <https://doi.org/10.1001/jama.2017.0686>. PMID: 28245334; PMCID: PMC5483855.

[28] Dabelea D, Mayer-Davis EJ, Saydah S, Imperatore G, Linder B, et al. (2014) SEARCH for Diabetes in Youth Study. Prevalence of type 1 and type 2 diabetes among children and adolescents from 2001 to 2009. *JAMA* **311**:1778-86. doi: <https://doi.org/10.1001/jama.2014.3201>. PMID: 24794371; PMCID: PMC4368900.

[29] Davids D, Gibson D, Johnson Q. (2016) Ethnobotanical survey of medicinal plants used to manage High Blood Pressure and Type 2 Diabetes Mellitus in Bitterfontein, Western Cape Province, South Africa. *J Ethnopharmacol* **194**:755-766. doi: <https://doi.org/10.1016/j.jep.2016.10.063>. PMID: 27780752.

[30] Giovannini P, Howes MJ, Edwards SE. (2016) Medicinal plants used in the traditional management of diabetes and its sequelae in Central America: A review. *J Ethnopharmacol* **184**:58-71. doi: <https://doi.org/10.1016/j.jep.2016.02.034>. PMID: 26924564.

[31] Astutik S, Pretzsch J, Ndzifon Kimengsi J. (2019) Asian Medicinal Plants' Production and Utilization Potentials: A Review. *Sustainability* **11**:5483. doi: <https://doi.org/10.3390/su11195483>.

[32] Oyebode O, Kandala NB, Chilton PJ, Lilford RJ. (2016) Use of traditional medicine in middle-income countries: a WHO-SAGE study. *Health Policy Plan* **31**:984-91. doi: <https://doi.org/10.1093/heropol/czw022>. PMID: 27033366; PMCID: PMC5013777.

[33] Safe S, Jayaraman A, Chapkin RS, Howard M, Mohankumar K, Shrestha R. (2021) Flavonoids: structure-function and mechanisms of action and opportunities for drug development. *Toxicol Res* **37**:147-162. doi: <https://doi.org/10.1007/s43188-020-00080-z>. PMID: 33868973; PMCID: PMC8007671.

[34] Beejmohun V, Peytavy-Izard M, Mignon C, Muscente-Paque D, Deplanque X, et al. (2014) Acute effect of Ceylon cinnamon extract on postprandial glycemia: alpha-amylase inhibition, starch tolerance test in rats, and randomized crossover clinical trial in healthy volunteers. *BMC Complement Altern Med* **14**:351. doi: <https://doi.org/10.1186/1472-6882-14-351>. PMID: 25249234; PMCID: PMC4246455.

[35] Salazar-García M, Corona JC. (2021) The Use of Natural Compounds as a Strategy to Counteract Oxidative Stress in Animal Models of Diabetes Mellitus. *Int J Mol Sci* **22**:7009. doi: <https://doi.org/10.3390/ijms22137009>.

[36] Djeridane A, Hamdi A, Bensania W, Cheffa K, Lakhdari I, et al. (2015) The in vitro evaluation of antioxidant activity, α -glucosidase and α -amylase enzyme inhibitory of natural phenolic extracts. *Diabetes Metab Syndr* **9**:324-31. doi: <https://doi.org/10.1016/j.dsx.2013.10.007>. PMID: 25470628.

[37] Kavishankar GB, Lakshmidhi N, Murthy SM, Prakash HS, Niranjana SR. (2011) Diabetes and medicinal plants-A review. *Int J Pharm Biomed Sci* **2**:65-80.

[38] Kamiya K, Hamabe W, Harada S, Murakami R, Tokuyama S, Satake T. (2008) Chemical constituents of *Morinda citrifolia* roots exhibit hypoglycemic effects in streptozotocin-induced diabetic mice. *Biol Pharm Bull* **31**:935-8. doi: <https://doi.org/10.1248/bpb.31.935>. PMID: 18451522.

[39] Horsfall A, Olabiyi O, Osinubi A, Noronha C, Okanlawon A. (2008) Antidiabetic effect of fruit juice of *Morinda Citrifolia* (Tahitian Noni Juices) on experimentally induced diabetic rats. *Nigerian J Health Biomedical Sci* **7**:34-37. doi: <https://doi.org/10.4314/njhbs.v7i2.11674>.

[40] Iwu MM, Igoboko OA, Okunji CO, Tempesta MS. (1990) Antidiabetic and aldose reductase activities of biflavonones of *Garcinia kola*. *J Pharm Pharmacol* **42**:290-2. doi: <https://doi.org/10.1111/j.1422-7158.1990.tb05412.x>. PMID: 1974302.

[41] Alade GO, Adebajo AC, Omobuwajo OR, Proksch P, Verspohl EJ. (2012) Quercetin, a minor constituent of the antihyperglycemic fraction of *Bauhinia monandra* leaf. *J Diabetes* **4**:439-41. doi: <https://doi.org/10.1111/j.1753-0407.2012.00222.x>. PMID: 22742491.

[42] Daisy P, Balasubramanian K, Rajalakshmi M, Eliza J, Selvaraj J. (2010) Insulin mimetic impact of Catechin isolated from *Cassia fistula* on the glucose oxidation and molecular mechanisms of glucose uptake on

Streptozotocin-induced diabetic Wistar rats. *Phytomedicine* **17**:28-36. doi: <https://doi.org/10.1016/j.phymed.2009.10.018>. PMID: 19931438.

[43] Koonyosing P, Uthaipibull C, Fucharoen S, Koumoutsea EV, Porter JB, Srichairatanakool S. (2019) Decrement in Cellular Iron and Reactive Oxygen Species, and Improvement of Insulin Secretion in a Pancreatic Cell Line Using Green Tea Extract. *Pancreas* **48**:636-643. doi: <https://doi.org/10.1097/MPA.00000000000001320>. PMID: 31091209; PMCID: PMC6553981.

[44] Wang JY, Nie YX, Dong BZ, Cai ZC, Zeng XK, et al. (2021) Quercetin protects islet β -cells from oxidation-induced apoptosis via Sirt3 in T2DM. *Iran J Basic Med Sci* **24**:629-635. doi: <https://doi.org/10.22038/ijbms.2021.52005.11792>. PMID: 34249264; PMCID: PMC8244598.

[45] Rahimi R, Nikfar S, Larijani B, Abdollahi M. (2005) A review on the role of antioxidants in the management of diabetes and its complications. *Biomed Pharmacother* **59**:365-73. doi: <https://doi.org/10.1016/j.biopha.2005.07.002>. PMID: 16081237.

[46] Li YG, Ji DF, Zhong S, Lv ZQ, Lin TB, et al. (2011) Hybrid of 1-deoxynojirimycin and polysaccharide from mulberry leaves treat diabetes mellitus by activating PDX-1/insulin-1 signaling pathway and regulating the expression of glucokinase, phosphoenolpyruvate carboxykinase and glucose-6-phosphatase in alloxan-induced diabetic mice. *J Ethnopharmacol* **134**:961-70. doi: <https://doi.org/10.1016/j.jep.2011.02.009>. PMID: 21333726.

[47] Akiyama S, Katsumata S, Suzuki K, Nakaya Y, Ishimi Y, Uehara M. (2009) Hypoglycemic and hypolipidemic effects of hesperidin and cyclodextrin-clathrated hesperetin in Goto-Kakizaki rats with type 2 diabetes. *Biosci Biotechnol Biochem* **73**:2779-82. doi: <https://doi.org/10.1271/bbb.90576>. PMID: 19966469.

[48] Hennebelle T, Weniger B, Joseph H, Sahpaz S, Bailleul F. (2009) Senna alata. *Fitoterapia* **80**:385-93. doi: <https://doi.org/10.1016/j.fitote.2009.05.008>. Epub 2009 May 14. PMID: 19446609.

[49] Campbell-Toft J, Mølgaard, Winther K. (2012) Harnessing the potential clinical use of medicinal plants as anti-diabetic agents. *Botanics: Targets and Therapy* **2**:7-19 doi: <https://doi.org/10.2147/BTAT.S17302>.

[50] Ezuruike UF, Prieto JM. (2014) The use of plants in the traditional management of diabetes in Nigeria: pharmacological and toxicological considerations. *J Ethnopharmacol* **155**:857-924. doi: <https://doi.org/10.1016/j.jep.2014.05.055>. PMID: 24929108.

[51] Aryal B, Raut BK, Bhattacharai S, Bhandari S, Tandan P, et al. (2022) Potential Therapeutic Applications of Plant-Derived Alkaloids against Inflammatory and Neurodegenerative Diseases. *Evid Based Complement Alternat Med* **2022**:7299778. doi: <https://doi.org/10.1155/2022/7299778>. PMID: 35310033; PMCID: PMC8926539.

[52] Odoh UE, Ezugwu CO. (2012) Anti-diabetic and toxicological studies of the alkaloids of Acanthus Montanus (Acanthaceae) leaf. *Planta Med* **78**: CL70. doi: <https://doi.org/10.1055/s-0032-1320305>

[53] Tiong SH, Looi CY, Hazni H, Arya A, Paydar M, et al. (2013) Antidiabetic and antioxidant properties of alkaloids from Catharanthus roseus (L.) G. Don. *Molecules* **18**:9770-84. doi: <https://doi.org/10.3390/molecules18089770>. PMID: 23955322; PMCID: PMC6270616.

[54] Agrawal R, Sethiya NK, Mishra SH. (2013) Antidiabetic activity of alkaloids of Aerva lanata roots on streptozotocin-nicotinamide induced type-II diabetes in rats. *Pharm Biol* **51**:635-42. doi: <https://doi.org/10.3109/13880209.2012.761244>. Epub 2013 Mar 26. PMID: 23527955.

[55] Jiang S, Chen C, Dong Q, Shao Y, Zhao X, et al. (2021) Alkaloids and phenolics identification in fruit of Nitraria tangutorum Bobr. by UPLC-Q-TOF-MS/MS and their α -glucosidase inhibitory effects in vivo and in vitro. *Food Chem* **364**:130412. doi: <https://doi.org/10.1016/j.foodchem.2021.130412>. Epub 2021 Jun 18. PMID: 34174646.

Author affiliation

1. Department Chemistry, College of Science, University of Baghdad, Baghdad, Iraq.

ORCID IDs:

1. Khalid A. Mohammed: <https://orcid.org/0009-0006-2092-0228>