Volume 12, Number 2: 26 - 31

ISSN: 2313-3937

2024

Review article

Exploring the Effects of Hydrogen Peroxide on Biofilm Development and Antibiotic Susceptibility

Hwazen Amer Shnyoor^{1,2}, Ayaid Khadem Zgair²

ABSTRACT

Hydrogen peroxide is one of the most important chemical compounds that affect the formation of biofilm of microorganisms and response to antimicrobial agents, and it can be used in war against pathogenic microorganisms in healthcare institutions. The current study aims to highlight the mechanism of the effect of H_2O_2 on biofilm formation, in addition to the possibility of using it to enhance the ability of antimicrobials to reduce the outbreak of infectious disease. In this review article, the impact of hydrogen peroxide on the susceptibility of pathogenic bacteria to antibiotics will be screened through the role of H_2O_2 in reducing biofilm formation. Here, the mechanism of the effect of H_2O_2 on the body of pathogenic bacteria through oxidation stress, eradication of biofilm, destruction of microbial membrane, and inhibition of microbial enzymes. The effect of hydrogen peroxide on biofilm formation depends on the concentration of hydrogen peroxide and the duration of exposure of microorganisms. At some times, low concentrations of H_2O_2 induce biofilm formation, and high concentrations may reduce the biofilm formation (and that may be dependent on the species of microorganism). The effect of H_2O_2 on biofilm formation is through reducing initial attachment as well as interfering with the maturation and maintenance of biofilm. Hydrogen peroxide has an important role in enhancing antibacterial agents, and this may contribute to redrawing the therapeutic strategy in the future. It can be concluded from the present study that H_2O_2 has a role in reducing the outbreak of infectious diseases and helps enhance the antibacterial war, which contributes to enhancing public health.

Keywords: Antibiotics, Bacterial susceptibility, Biofilm, Developing antibacterial agents, Hydrogen peroxide.

Citation: Shnyoor HA, Zgair AK. (2024) Exploring the Effects of Hydrogen Peroxide on Biofilm Development and Antibiotic Susceptibility. *World J Exp Biosci* **12**: 26 - 31.

Received: July 11, 2024; Revised: August 22, 2024; Accepted: August 27, 2024; Published: September 19.

1. INTRODUCTION

Biofilms are complex microbial communities that form and attach to different surfaces encased in a self-produced extracellular matrix [1]. The biofilm is not just found in plumbing but is also in all sorts of natural and industrial settings. They can form on various surfaces and are often found in nutrient-rich environments. In many instances, processes within natural biofilms are analogous to what happens in artificial or industrial setups [2]. Previous studies showed the elimination of bacterial biomass and biofilm by using hydrogen peroxide (H₂O₂) [3]. Most biofilms are found in environments created by humans: inside houses, in industry, and the medical devices that produce

the defects [2]. While it is expected that H_2O_2 would effectively eradicate biofilms by penetrating and disrupting the microbial cells within, recent studies indicate that its effects are more nuanced [4]. Research using computational modeling has demonstrated that biofilms can exhibit unexpected behaviors when exposed to hydrogen peroxide. Previous studies showed that a particular concentration of H2O2 may not eradicate the biofilm but increase the thickness of the biofilm [5]. The dead biomass in the biofilm of bacterial culture protects the viable bacterial cells from the effect of H_2O_2 by neutralizing its effect that protects viable cells in the deep area in the biofilm from oxi-

Copyright: © Hwazen A. Shnyoor & Ayaid K, Zgair. This is an open-access article distributed under the terms of the Creative Commons Attribution. International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

^{*} Correspondence: **Dr. Ayaid Khadem Zgair.** E. mail: ayaid.zgair@sc.uobaghdad.edu.iq
Department of Biology, College of Science, University of Baghdad, Baghdad, Iraq.
Full list of author information is available at the end of the article.

oxi stress that yields from the H2O2 [6]. This finding creates the changes to using the disinfected agents against the bacterial biomass in the hospitals that lead to re-evaluating the strategies of using the oxidative agents as disincentive agents on the floor of the health institutions.

The relationship between hydrogen peroxide and antibiotic effectiveness is a key aspect of this review article. When present, H_2O_2 can change how well antibiotics work against biofilms. Biofilms consist of bacteria that stick to surfaces, and they can form in human bodies, for instance, on indwelling catheters. There's currently no good way to treat such infections. The using of H_2O_2 and antibiotics together may reflect positively on treating the bacterial infection and ultimately produce good patient health [7].

Several factors interfere with the effect effects of H2O2 on biofilm development which requires investigation when the researchers want to understand biofilms and their susceptibility to antibiotics. In this review, the effect of H2O2 on the dynamics of biofilm formation to build strategies for managing the bacterial infections that are related to biofilm formation. This will help to improve the effectiveness of antibiotics against the bacterial infection associated with biofilm formation.

2. OVERVIEW of BIOFILM

Biofilm is a biomass of microorganisms that communicate together at the biotic or abiotic surfaces. Commonly known as polysaccharides, proteins, and nucleic acids [1]. This matrix maintains structural integrity and protects biofilms, which enables them to exist in a variety of environments, from ecosystems to medical devices [2]. The development of biofilms occurs in several stages: initial attachment of planktonic cells to a surface, irreversible adhesion due to the production of extrace-Ilular polymeric substances, growth into three-dimensional structures that are quite complex, and the shedding of cells that can settle in a new region [8]. Infections are particularly difficult to treat because biofilms are refractory to the host immune response as well as antimicrobial agents [9]. The bacteria within the biofilm communicate with one another through the mechanism of bacterial quorum sensing. This phenomenon is a cell-tocell communication that uses gene expression controlled by the population density of the microorganisms. Such communication also enables synchronized actions such as nutrient acquisition and biofilm development [10]. Biofilms can be useful in cases of bioremediation and wastewater treatment, however, in the field of medicine, they are of great concern due to their role in chronic infections and device-associated infections [11]. The understanding of biofilm development and related resistance mechanisms to antibiotics is highly important in controlling the infections that are related to biofilm formation

2.1. Important of antibiotic Susceptibility

Antibiotic resistance in bacteria that form biofilm strongly has become a focus of attention because biofilm poses specific problems in both clinical and environmental settings. Biofilms are structured communities of bacteria locked in an extracellular matrix and there is often a marked difference in antibiotic susceptibility between biofilms and planktonic free-floating [12] microbes. The majority of bacteria in biofilms acquire higher resistance to most classes of antibiotics. The basis of these resistances includes factors like restricted diffusion of the antimicrobial agents, and differentiation of cells in the biofilm

with slow growth rates reason these cells would be less affected by antibiotics and these cells could also produce enzymes able to inactivate the antimicrobial agents. For instance, Staphylococcus aureus biofilm-forming strains have been reported to have high levels of resistance to beta-lactams which pose a critical challenge in the clinical context of biofilm-mediated infections [13]. Biofilms are involved in about 65-80 percent of chronic infections and also in device-associated and implant-associated infections. The ability of biofilm-forming bacteria to escape most of the conventional antibiotics treatment makes it easier to deal with the infection however such patients may need to go through black box procedures including explanation of infected devices of the patients [14].

It is important to gain insight into the association between biofilm formation and antimicrobial sensitivity for efficient treatment methods to be adopted. According to research, some specific antibiotics can stimulate the establishment of biofilms under low doses, pointing toward a more complex relationship of biofilms with antibiotics [15]. With this understanding, the development of other novel means that may help to target biofilms or enhance the effectiveness of already available antibiotics can be achieved. The appearance of multi-drug resistance (MDR) and extensively drug-resistant (XDR) biofilm strains creates a burden in the community. Regular evaluation and monitoring of biofilm-forming bacteria for antibiotic resistance is essential to prevent and treat infections appropriately [16]. Indeed, the investigation of antibiotic resistance in biofilm-forming bacteria is relevant for overcoming the threats of persistent infections and creating potential therapeutic solutions. Ultimately, knowing these phenomena will be valuable in improving clinical outcomes and the management of biofilm infectious diseases.

3. MECHANISMS of ACTION of HYDROGEN PEROXIDE

 $\rm H_2O_2$ is an antimicrobial agent that is widely used for its disinfectant equipment, especially medical devices. Its mechanism of action consists of several biochemical processes that contribute to its effectiveness against many microorganisms including bacteria, viruses, and fungi

- Production of reactive oxygen species (ROS): H₂O₂ produces reactive oxygen species. Especially hydroxyl radicals are highly reactive and destroy cellular components, lipids, ROS, proteins, nucleic acids, etc. They can oxidize important biomolecules. This leads to cell death. Oxidative damage disrupts cell membranes and impairs vital cellular functions. Ultimately, it leads to bacterial fragmentation and death [17].
- Membrane Damage: H₂O₂ penetrates microbial cell walls and membranes, leading to increased permeability. This disruption allows for the leakage of intracellular contents and the loss of essential ions, further compromising cell integrity. The damage to membrane lipids initiates further oxidative stress within the cell [17,18].
- Inhibition of Enzymatic Activity: The H2O2 can inhibit the important metabolic enzymes that are involved in bacterial metabolism that happen by changing the enzyme structure and kinetics process through the oxidation process that may reflect on the bacteria metabolism process that is critical in bacterial growth and survival [19].
- Biofilm Disruption: Prior research suggests that H₂O₂ can
 effectively reduce the biofilm biomass of different species of
 bacteria. It disturbs the extracellular polymeric substances
 (EPS) which are the biofilm matrix that protects embedded
 bacteria from exposure to antimicrobial agents. Hydrogen

peroxide has been shown to dramatically reduce biofilm density and viability with relatively short exposure times [18,20].

Hydrogen peroxide as an antimicrobial agent and its antibiotic mechanisms of action emphasize the efficacy of hydrogen peroxide on various microbial species through oxidative damage, disruption to membranes, and biofilm minimization. Recognizing these mechanisms is crucial for maximizing its potential with applications such as in health care where infectious agents must be contained.

4. IMPACT of H₂O₂ on BIOFILM FORMATION

The previous studies on the effect of Hydrogen peroxide have shown two sward edges. In some cases, the H2O2 promotes biofilm formation and in other studies, the H2O2 eradicates the biofilm biomass. That is dependent on the bacteria population microenvironments and the concentration of H2O2 in the environments of bacterial growth [21]. Duan et al (2016) reported that H2O2 promotes *Aggregatibacter actinomycetemcomitans* and *Streptococcus parasanguinis* to form biofilm [21]. Other investigators reported that H2O2 decreases and eradicates the biofilm formation [22]. There are several keys to the effect of H2O2 on the biofilm formation.

- Anti-Biofilm Activity: H₂O₂ is effective in eradicating by biofilms formed various bacterial species. including Escherichia coli. Research indicates concentrations of H₂O₂ at or above 6.25% (v/v), when applied at elevated temperatures (≥40 °C) for sufficient exposure times (≥25 min), can significantly reduce biofilm biomass and viability. This suggests that H₂O₂ disrupts the structural integrity of biofilms, leading to cell lysis and death [23].
- Temperature and Concentration of H2O2: The
 effectiveness of H₂O₂ against biofilms is influenced by
 factors such as concentration, temperature, and contact
 time. Higher concentrations and temperatures enhance its
 ability to penetrate the biofilm matrix and exert lethal effects
 on embedded bacteria [23].
- Raise of Biofilm Formation: Exogenous addition of H₂O₂ in low concentration promotes the production of biofilm in certain bacterial isolates, i.e. Acinetobacter oleivorans. This happens because the oxidative stress responses enhance the production of extracellular polymeric substances (EPS), which are vital in biofilm development. The other researchers found that H₂O₂ in certain concentrations stimulates genes associated with EPS production, thereby facilitating initial biofilm formation [24].
- Tolerance Mechanisms: The biomass of biofilms shows tolerance to H₂O₂ because of the neutralization of the compound by dead biomass of bacteria. This will help viable cells in the biofilm to survive and get rid of the oxidative stress of H₂O₂. Studies have shown that prolonged exposure to sub-lethal concentrations of H₂O₂ leads to increased thickness and resilience of biofilms in particular bacteria species [6].
- Long-Term Effects on Biofilms: Treatment with H₂O₂ concentrations (low concentrations) fails to eradicate established biofilms will stimulate bacterial biomass to regrowth after treatment with H2O2. Cyanobacteria biofilms exposed to H₂O₂ demonstrated increased biomass two weeks after treatment, indicating that disinfecting agents might inadvertently promote biofilm resilience if not applied in sufficient concentration [25].

4.1. H₂O₂ inhibits the initial attachment

Hydrogen Peroxide (H₂O₂) has a major effect on the initial attachment of bacteria which is, an important step during biofilm formation. The influence of H2O2 on this process could be elucidated by a variety of mechanisms. Initial Inhibition of Attachment, hydrogen peroxide inhibits the ability of bacteria to initially attach to the surface via producing various ROS species experienced by bacterial oxidative stress which can break down cell membranes thus hampering the attachment of bacteria. This is especially important in settings where biofilm formation is considered to be detrimental, such as medical devices and water systems [26]. Concentration-Dependent Effects, the effect of H2O2 on bacterial adhesion is concentrationdependent. Hydrogen Peroxide at higher concentrations (≥6.25% v/v) has previously demonstrated the ability to eradicate preformed biofilms and inhibit subsequent biofilm development, resulting in more than a 3-log reduction in total biofilm formation [14]. In comparison, lower concentrations may not have adequate antimicrobial activity to prevent attachment [23, 24]. Oxidative Stress Response: Exposure H₂O₂ then bacteria activate a stress response mechanism that temporarily inhibits bacterial adhesion to abiotic surfaces which the effect is the first step in biofilm formation. The oxidative stress induces the expression of detoxification and damage repair genes (arcA, mdoG, and tus) in E. coli [27], which can change the bacteria's properties affecting its adhesion Exopolysaccharides (EPS) while in other contexts H2O2 can inhibit initial attachment and adhesion of bacteria, it has been observed that biofilm formation is promoted following H2O2 treatment due to the induction of exopolysaccharides (EPS) production from some species like A. oleivorans. This dual role emphasizes the paradoxical nature of H₂O₂ effects when lower concentration can contribute to EPS production under certain conditions and favor later steps of biofilm development [24]. Competitive Dynamics: In competitive environments, some species produce H₂O₂ inhibits the growth of non-adapted ones and favors the growth of adapted species instead. The developing biofilm community composition can be shaped by such selective pressures [26]. Therefore, the antimicrobial action and oxidative stress-inducing capability of H₂O₂ play an important role in the meditating effect maintained on bacterial initial attachment and biofilm formation. It mostly has an inhibitory effect on initial attachment at higher concentrations but also increases biofilm formation generally by inducing exopolysaccharides (EPS) production [26]. Log in or register to get access to the full content.

4.2. Interference with Maturation and Maintenance

The impact of H2O2 on the dynamics of biofilm development, maturation, and maintenance is complex. Hydrogen peroxide influences the stability and maintenance of biofilm. The concentration, duration of exposure, and type of microbial community significantly influence the effect of hydrogen peroxide on biofilm development.

4.2.1. Inhibition of Biofilm Maturation

 H₂O₂ induces oxidative stress: Microbial cells subjected to elevated levels of hydrogen peroxide and H2O2 will undergo oxidative stress, hence impeding biofilm development. Hydrogen peroxide can impede bacterial proliferation by destroying biological constituents such as membranes and DNA, therefore breaking stable biofilm formations [18]. The prior work demonstrated that H_2O_2 effectively reduces biofilm thickness and the survival of *Staphylococcus epidermidis*, with optimal concentrations ranging from 3% to 5% [18].

Disruption of Extracellular Polymeric Substances (EPS):
 H₂O₂ affects the production and integrity of EPS, which are crucial for biofilm structure. The oxidative action can degrade EPS components, leading to a less stable biofilm matrix and facilitating detachment of Streptococcus mutans bacterial cells from the biofilm and also killing the bacteria and helping in the maintenance of the patient teeth [28].

4.2.2. Impact on Biofilm Maintenance

- Catalase Activity: The presence of catalase-producing bacteria within a biofilm neutralizes H₂O₂, allowing for the survival of deeper layers while protecting them from oxidative damage. This ability to neutralize H₂O₂ contributes to the maintenance of biofilms that were formed by *Pseudomonas aeruginosa* even during continuous exposure to the disinfectant. Studies using computational models have shown that dead biomass within a biofilm can absorb H₂O₂, providing a protective barrier for living cells deeper within the structure [29].
- Concentration Thresholds: Research indicates that sustained low concentrations of H₂O₂ may not be effective in completely eradicating biofilms. Instead, treatments need to exceed a critical threshold concentration to effectively disrupt biofilm maintenance mechanisms. Continuous low-dose exposure may allow for adaptive responses that enhance biofilm resilience. The previous studies support that high concentrations of H₂O₂ can eradicate the biofilm formed by *E. coli* [23] while low concentrations of endogenous H₂O₂ increase the EPS produced by *A. oleivorans* and enhance biofilm formation [24].

4.2.3. Adaptive Responses

Non-lethal exposure to H_2O_2 induces a priming response in bacteria, enhancing their survival against subsequent lethal doses. This adaptive response may involve the up-regulation of stress response pathways and increased efficiency in removing H_2O_2 from their environment, ultimately contributing to biofilm resilience [27].

Hydrogen peroxide significantly interferes with both the maturation and maintenance of biofilms through oxidative stress, disruption of EPS integrity, and the influence of microbial adaptive responses. While it can effectively inhibit biofilm development at higher concentrations, its interaction with catalase activity and potential priming effects complicate its use as a disinfectant.

5. H₂O₂ as an ADJUNCT to ANTIBIOTIC THERAPY

In recent years an interest in the application use of H_2O_2 as a potential adjunct to antibiotic therapy, particularly in the context of combating antibiotic-resistant infections and enhancing the efficacy of existing antimicrobial treatments. Its mechanisms of action are dependent on the oxidation damage of the bacterial cells, biofilm disruption, and synergistic effect with other antimicrobial agents [30]. There wide spectrum of applications of using hydrogen peroxide in clinical applications. Recent

advancements have shown that H_2O_2 when combined with blue light therapy, can effectively kill MRSA by destabilizing cell membranes and allowing H_2O_2 to penetrate bacterial cells. This combination has been reported to achieve up to 99.9% bacterial reduction [31]. H_2O_2 has been explored in various chronic infection models. For instance, its application in dental unit water systems has shown significant reductions in microbial load, indicating its potential for broader clinical applications [30].

Despite its potential benefits, there are limitations to the use of $\rm H_2O_2$ as an adjunct therapy. The effectiveness of $\rm H_2O_2$ can vary based on concentration and exposure duration. High concentrations may be necessary for effective antimicrobial action but could also lead to tissue damage if not carefully managed [23,24]. The presence of catalase-producing bacteria can neutralize $\rm H_2O_2$, reducing its effectiveness in certain environments, such as within biofilms or in the bloodstream [32]. $\rm H_2O_2$ may enhance antibiotic efficacy, but there is a concern regarding potential resistance development in bacteria exposed to sub-lethal concentrations over time [27].

Hydrogen peroxide shows promise as an adjunct to antibiotic therapy by disrupting biofilms, enhancing antibiotic penetration, and exerting oxidative stress on bacteria. Its application in treating resistant infections like MRSA highlights its potential clinical utility. However, careful consideration of concentration, exposure time, and microbial characteristics is essential for optimizing its use. Further research is needed to establish standardized protocols for combining H_2O_2 with antibiotics in clinical settings to maximize therapeutic outcomes while minimizing risks.

6.1. Synergistic Effects with Antibiotics

Hydrogen peroxide's synergistic effects when combined with various antibiotics and other antimicrobial agents illustrate its potential as a treatment for bacterial infections caused by highresistance bacteria. The use of H₂O₂ in combination with these standard treatments can enhance their antibacterial effects in several ways, including by causing oxidative stress, which makes the bacteria much more susceptible to the simultaneously applied antibiotics; at high enough concentrations, H2O2 itself can do this as can other standard treatments like chlorhexidine. Feuerstein et al. (2006) directed another line of research that was aimed at discovering the exact nature of the "synergistic" effect that they had previously reported. They found that blue light and H₂O₂ together exhibited over 96% growth inhibition against Streptococcus mutans compared to the much lower growth inhibition levels when these agents were applied separately. The investigators hypothesized that the use of blue light to activate H₂O₂ resulted in a quantum leap in the photosensitization potential of the H₂O₂ and was responsible for the production of a much larger quantity of the reactive oxygen species (ROS) that they had previously identified as being responsible for the enhanced antibacterial effect produced by the use of these two agents in concert [33]. The antibacterial potency of the combination of chlorhexidine (CHX) and hydrogen peroxide (H2O2) was equivalent to that of sodium hypochlorite (NaOCI). Thus, it could serve as a useful root canal irrigant. The combined antibacterial effect of CHX and H₂O₂ is greater than either agent could accomplish on its own. The H₂O₂ not only penetrated the bacterial cell walls better than NaOCI could, but it also worked with the CHX to kill off the resistant cells that either agent wouldn't have been able to vanquish alone [34]. Alkawareek et al. (2019) found that silver nanoparticles also work synergistically with H2O2 to kill off bacteria [35]. This synergy came from a Fenton-like reaction where AgNP and H₂O₂ interacted. They produced highly reactive hydroxyl radicals [35]. This was good; these radicals enhanced the antibacterial activity we were studying. There was a downside, though. If not managed properly, that much reactivity could lead to toxicity. So, we managed it. We moderated the interaction of H₂O₂ with gluconic acid in honey. Then we hit the target. We got enhanced antibacterial activity against E. coli without too much toxicity [36]. The synergy between cold atmospheric plasma and hydrogen peroxide was revealed to possess powerful and unique bacterial disinfection properties in a recent study. When applied to notoriously difficult-to-kill bacteria like Enterococcus faecalis, combination of cold plasma and H₂O₂ got the job done and left tissues (the target of the treatment) in a state that was conducive to healing. Dramatic improvements were noted in the efficiency of this plasma plus H2O2 approach compared with using either component alone [37].

6. CONCLUSION

Hydrogen peroxide plays a certain role in biofilm formation as an antimicrobial agent and impacts biofilm formation. Its ability to disrupt biofilm biomass is mediated by inducing oxidative stress. Role of H2O2 in eradicating the biofilm by preventing the initial adhesion of bacteria and maturation of biofilm. The role of H2O2 in biofilm production and bacterial resistance to antibiotics call for the redesign the treatment strategies. The different concentrations and duration of exposure of H2O2, the species of microorganism play a central role of H2O2 in biofilm formation. The synergistic effect of H2O2 and antibiotic increase the antibacterial effect of antibiotic and help in improving the antibacterial effect of antimicrobial agents. Emphasizing collaboration between microbiologists, healthcare practitioners, and researchers will pave the way for innovative and effective solutions to the ongoing problem of biofilm-associated infections.

Acknowledgments

At this stage, I would like to thank the staff of all hospitals in Babylon provinces and the Department of Biology, College of Science, University of Baghdad, for their valuable assistance in preparing the review article.

Funding information

This work received no specific grant from any funding agency.

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical Approval

This review was approved by the Scientific Committee of the University of Baghdad, Baghdad, Iraq (1415; 16-04-2024).

Author contributions

Hwazen A. Shnyoor: Investigation; Methodology; Project administration; Resources; Supervision; Validation; Roles/Writing - original draft; and Writing - review & editing.

Ayaid K. Zgair. Conceptualization, Data curation, and Formal analysis, Roles/Writing - original draft; Visualization and Writing - review & editing.

6. REFERENCES

[1] Karygianni L, Ren Z, Koo H, Thurnheer T. (2020) Biofilm Matrixome: Extracellular Components in Structured Microbial Communities. Trends Microbiol 28:668-681. doi: 10.1016/j.tim.2020.03.016. Epub 2020 Apr 21. PMID: 32663461.

- [2] Pandit A, Adholeya A, Cahill D, Brau L, Kochar M. (2020) Microbial biofilms in nature: unlocking their potential for agricultural applications. J Appl Microbiol 129:199-211. doi: 10.1111/jam.14609. Epub 2020 Feb 20. PMID: 32034822.
- [3] Preethi, Rajesh Banu J, Kavitha S, Yukesh Kannah R, Varjani S, Gunasekaran M. (2022) Mild hydrogen peroxide interceded bacterial disintegration of waste activated sludge for efficient biomethane production. Sci Total Environ 817:152873. doi: 10.1016/j.scitotenv.2021.152873. Epub 2022 Jan 5. PMID: 34998769.
- [4] Song J, Hong L, Zou X, Alshawwa H, Zhao Y, et al. (2022) A Self-Supplying H₂O₂ Modified Nanozyme-Loaded Hydrogel for Root Canal Biofilm Eradication. *Int J Mol Sci* 23:10107. doi: 10.3390/ijms231710107. PMID: 36077503; PMCID: PMC9456354.
- [5] Li M, Lan X, Han X, Shi S, Sun H, et al. (2021) Acid-Induced Self-Catalyzing Platform Based on Dextran-Coated Copper Peroxide Nanoaggregates for Biofilm Treatment. ACS Appl Mater Interfaces 13:29269-29280. doi: 10.1021/acsami.1c03409. Epub 2021 Jun 18. PMID: 34143595.
- [6] Stewart PS, Owkes M. (2023) Simulation of catalase-dependent tolerance of microbial biofilm to hydrogen peroxide with a biofilm computer model. NPJ Biofilms Microbiomes 9:60. doi: 10.1038/s41522-023-00426-z. PMID: 37612330; PMCID: PMC10447567.
- [7] Vatansever F, de Melo WC, Avci P, Vecchio D, Sadasivam M, et al. (2013) Antimicrobial strategies centered around reactive oxygen species--bactericidal antibiotics, photodynamic therapy, and beyond. FEMS Microbiol Rev 37:955-89. doi: 10.1111/1574-6976.12026. Epub 2013 Jul 25. PMID: 23802986; PMCID: PMC3791156.
- [8] Di Martino P. (2018) Extracellular polymeric substances, a key element in understanding biofilm phenotype. AIMS Microbiol 4:274-288. doi: 10.3934/microbiol.2018.2.274. PMID: 31294215; PMCID: PMC6604936.
- [9] Macià MD, Rojo-Molinero E, Oliver A. (2014) Antimicrobial susceptibility testing in biofilm-growing bacteria. *Clin Microbiol Infect* 20:981-90. doi: 10.1111/1469-0691.12651. Epub 2014 Jun 14. PMID: 24766583.
- [10] Prazdnova EV, Gorovtsov AV, Vasilchenko NG, Kulikov MP, Statsenko VN, et al. (2022) Quorum-Sensing Inhibition by Gram-Positive Bacteria. Microorganisms 10:350. doi: 10.3390/microorganisms10020350. PMID: 35208805; PMCID: PMC8875677.
- [11] Mirghani R, Saba T, Khaliq H, Mitchell J, Do L, et al. (2022) Biofilms: Formation, drug resistance and alternatives to conventional approaches. AIMS Microbiol 8:239-277. doi: 10.3934/microbiol.2022019. PMID: 36317001; PMCID: PMC9576500.
- [12] Han Y, Jiang N, Xu H, Yuan Z, Xiu J, et al. (2023) Extracellular Matrix Rigidities Regulate the Tricarboxylic Acid Cycle and Antibiotic Resistance of Three-Dimensionally Confined Bacterial Microcolonies. Adv Sci (Weinh) 10:e2206153. doi: 10.1002/advs.202206153. Epub 2023 Jan 19. PMID: 36658695; PMCID: PMC10037996.
- [13] Pajohesh R, Tajbakhsh E, Momtaz H, Rahimi E. (2022) Relationship between Biofilm Formation and Antibiotic Resistance and Adherence Genes in Staphylococcus aureus Strains Isolated from Raw Cow Milk in Shahrekord, Iran. Int J Microbiol 2022:6435774. doi: 10.1155/2022/6435774. PMID: 36329896; PMCID: PMC9626243.
- [14] Grooters KE, Ku JC, Richter DM, Krinock MJ, Minor A, et al. (2024) Strategies for combating antibiotic resistance in bacterial biofilms. Front Cell Infect Microbiol 14:1352273. doi: 10.3389/fcimb.2024.1352273.
- [15] Qian W, Li X, Yang M, Liu C, Kong Y, et al. (2022) Relationship Between Antibiotic Resistance, Biofilm Formation, and Biofilm-Specific Resistance in Escherichia coli Isolates from Ningbo, China. Infect Drug Resist 15:2865-2878. doi: 10.2147/IDR.S363652. PMID: 35686192; PMCID: PMC9172925.
- [16] Blanco-Cabra N, López-Martínez MJ, Arévalo-Jaimes BV, Martin-Gómez MT, Samitier J, Torrents E. (2021) A new BiofilmChip device for testing biofilm formation and antibiotic susceptibility. NPJ Biofilms Microbiomes 7:62. doi: 10.1038/s41522-021-00236-1. PMID: 34344902: PMCID: PMC8333102.
- [17] Shahriari S, Mohammadi Z, Mokhtari MM, Yousefi R. (2011) Effect of hydrogen peroxide on the antibacterial substantivity of chlorhexidine. Int J Dent 2010:946384. doi: 10.1155/2010/946384. Epub 2011 Jan 19. PMID: 21318180; PMCID: PMC3034917.

- [18] Presterl E, Suchomel M, Eder M, Reichmann S, Lassnigg A, et al. (2007) Effects of alcohols, povidone-iodine and hydrogen peroxide on biofilms of Staphylococcus epidermidis. J Antimicrob Chemother 60:417-20. doi: 10.1093/jac/dkm221. Epub 2007 Jun 22. PMID: 17586808.
- [19] Juven BJ, Pierson MD. (1996) Antibacterial Effects of Hydrogen Peroxide and Methods for Its Detection and Quantitation. J Food Prot 59:1233-1241. doi: 10.4315/0362-028X-59.11.1233. PMID: 31195444.
- [20] Lineback CB, Nkemngong CA, Wu ST, Li X, Teska PJ, Oliver HF. (2018) Hydrogen peroxide and sodium hypochlorite disinfectants are more effective against Staphylococcus aureus and Pseudomonas aeruginosa biofilms than quaternary ammonium compounds. Antimicrob Resist Infect Control 7:154. doi: 10.1186/s13756-018-0447-5. PMID: 30568790; PMCID: PMC6298007.
- [21] Duan D, Scoffield JA, Zhou X, Wu H. (2016) Fine-tuned production of hydrogen peroxide promotes biofilm formation of Streptococcus parasanguinis by a pathogenic cohabitant Aggregatibacter actinomycetemcomitans. Environ Microbiol 18:4023-4036. doi: 10.1111/1462-2920.13425. Epub 2016 Jul 22. PMID: 27348605; PMCID: PMC5118171.
- [22] Perumal PK, Wand ME, Sutton JM, Bock LJ. (2014) Evaluation of the effectiveness of hydrogen-peroxide-based disinfectants on biofilms formed by Gram-negative pathogens. J Hosp Infect 87:227-33. doi: 10.1016/j.jhin.2014.05.004. Epub 2014 Jun 5. PMID: 24957804.
- [23] Farjami A, Jalilzadeh S, Siahi-Shadbad M, Lotfipour F. (2022) The anti-biofilm activity of hydrogen peroxide against *Escherichia coli* strain FL-Tbz isolated from a pharmaceutical water system. *J Water Health* 20:1497-1505. doi: 10.2166/wh.2022.061. PMID: 36308494.
- [24] Jang IA, Kim J, Park W. (2016) Endogenous hydrogen peroxide increases biofilm formation by inducing exopolysaccharide production in *Acinetobacter oleivorans* DR1. Sci Rep 6:21121. doi: 10.1038/srep21121. PMID: 26884212; PMCID: PMC4756669.
- [25] Romeu MJ, Morais J, Vasconcelos V, Mergulhão F. (2023) Effect of Hydrogen Peroxide on Cyanobacterial Biofilms. Antibiotics (Basel) 12:1450. doi: 10.3390/antibiotics12091450. PMID: 37760746; PMCID: PMC10525773.
- [26] Zhu L, Kreth J. (2012) The role of hydrogen peroxide in environmental adaptation of oral microbial communities. Oxid Med Cell Longev 2012:717843. doi: 10.1155/2012/717843. Epub 2012 Jul 16. PMID: 22848782; PMCID: PMC3405655.
- [27] Rodríguez-Rojas A, Kim JJ, Johnston PR, Makarova O, Eravci M, et al. (2020) Non-lethal exposure to H202 boosts bacterial survival and evolvability against oxidative stress. PLoS Genet 16:e1008649. doi: 10.1371/journal.pgen.1008649. PMID: 32163413; PMCID: PMC7093028.
 - Author affiliation
 - 1. Ministry of health, Babylon health direct Alhashmia-hospital.
 - 2. Department of Biology, College of Science, University of Baghdad

ORCID IDs:

Ayaid K. Zgair: https://orcid.org/my-orcid?orcid=0000-0002-2356-3338

- [28] Ren Z, Kim D, Paula AJ, Hwang G, Liu Y, et al. (2019) Dual-Targeting Approach Degrades Biofilm Matrix and Enhances Bacterial Killing. J Dent Res 98:322-330. doi: 10.1177/0022034518818480. Epub 2019 Jan 24. PMID: 30678538.
- [29] Stewart PS, Roe F, Rayner J, Elkins JG, Lewandowski Z, et al. (2000) Effect of catalase on hydrogen peroxide penetration into Pseudomonas aeruginosa biofilms. Appl Environ Microbiol 66:836-8. doi: 10.1128/AEM.66.2.836-838.2000. PMID: 10653761; PMCID: PMC91906.
- [30] Hoogenkamp MA, Mazurel D, Deutekom-Mulder E, de Soet JJ. (2023) The consistent application of hydrogen peroxide controls biofilm growth and removes Vermamoeba vermiformis from multikingdom in-vitro dental unit water biofilms. Biofilm 5:100132. doi: 10.1016/j.biofilm.2023.100132. PMID: 37346320; PMCID: PMC10279787.
- [31] Dong PT, Mohammad H, Hui J, Leanse LG, Li J, et al. (2019) Photolysis of Staphyloxanthin in Methicillin-Resistant Staphylococcus aureus Potentiates Killing by Reactive Oxygen Species. Adv Sci (Weinh) 6:1900030. doi: 10.1002/advs.201900030. PMID: 31179216; PMCID: PMC6548961.
- [32] Shenep JL, Stokes DC, Hughes WT. (1985) Lack of antibacterial activity after intravenous hydrogen peroxide infusion in experimental Escherichia coli sepsis. Infect Immun 48:607-10. doi: 10.1128/iai.48.3.607-610.1985. PMID: 3888840; PMCID: PMC261202.
- [33] Feuerstein O, Moreinos D, Steinberg D. (2006) Synergic antibacterial effect between visible light and hydrogen peroxide on Streptococcus mutans. J Antimicrob Chemother 57:872-6. doi: 10.1093/jac/dkl070. Epub 2006 Mar 13. PMID: 16533827.
- [34] Hasheminia S, Farhad AR, Saatchi M, Rajabzadeh M. (2013) Synergistic antibacterial activity of chlorhexidine and hydrogen peroxide against *Enterococcus faecalis*. J Oral Sci 55:275-80. doi: 10.2334/josnusd.55.275. PMID: 24351914.
- [35] Alkawareek MY, Bahlool A, Abulateefeh SR, Alkilany AM. (2019) Synergistic antibacterial activity of silver nanoparticles and hydrogen peroxide. PLoS One 14:e0220575. doi: 10.1371/journal.pone.0220575. PMID: 31393906; PMCID: PMC6687290.
- [36] Masoura M, Passaretti P, Overton TW, Lund PA, Gkatzionis K. (2020) Use of a model to understand the synergies underlying the antibacterial mechanism of H2O2-producing honeys. Sci Rep 10:17692. doi: 10.1038/s41598-020-74937-6. PMID: 33077785; PMCID: PMC7573686.
- [37] Baik KY, Jo H, Ki SH, Kwon GC, Cho G. (2023) Synergistic effect of hydrogen peroxide and cold atmospheric pressure plasma-jet for microbial disinfection. *Appl Sci* 13: 3324. https://doi.org/10.3390/app13053324